

**THIS PAGE BLANK (USPTO)**

Exhibit J

L3 ANSWER 16 OF 21 MEDLINE DUPLICATE 14  
 ACCESSION NUMBER: 1998307272 MEDLINE  
 DOCUMENT NUMBER: 98307272 PubMed ID: 9645365  
 TITLE: Identification of a human member of the Ly-49 multigene family.  
 AUTHOR: Westgaard I H; Berg S F; Orstavik S; Fossum S; Dissen E  
 CORPORATE SOURCE: Department of Anatomy, Institute of Basic Medical Sciences,  
 University of Oslo, Norway.. i.h.westgaard@basalmed.uio.no  
 SOURCE: EUROPEAN JOURNAL OF IMMUNOLOGY, (1998 Jun) 28 (6) 1839-46.  
 Journal code: 1273201. ISSN: 0014-2980.  
 PUB. COUNTRY: GERMANY: Germany, Federal Republic of  
 Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 OTHER SOURCE: GENBANK-AF047445  
 ENTRY MONTH: 199807  
 ENTRY DATE: Entered STN: 19980723  
 Last Updated on STN: 20000303  
 Entered Medline: 19980715  
 AB Three classes of multigene family-encoded receptors enable NK cells to discriminate between polymorphic MHC class I molecules: Ly-49 homodimers, CD94/NKG2 heterodimers and the killer cell inhibitory receptors (KIR). Of these, CD94/NKG2 has been characterized in both rodents and humans. In contrast, Ly-49 family members have hitherto been found only in rodents, and KIR molecules only in the human. In this report, we describe a human cDNA, termed Ly-49L, that constitutes the first human member of the Ly-49 multi-gene family. Compared with rodent Ly-49 molecules, the Ly-49L sequence contains a premature stop codon and predicts a truncated protein that lacks the distal part of a C-terminal **lectin** domain. Evidence is presented that the premature stop codon results from incomplete excision of the intron between the first two **lectin** domain exons. **Splice variants** predicting a full-size Ly-49L protein were not detected. As demonstrated by Northern blot analysis, Ly-49L was transcribed by IL-2-activated NK cells, but not by freshly isolated B or T cells. PCR screening of a 22-clone yeast artificial chromosome contig localized the LY49L locus to the human NK gene complex on chromosome 12p12-p13. Southern blot analysis of genomic DNA showed a simple pattern with a full-length Ly-49L probe at low stringency hybridization conditions, suggesting that Ly-49L may be the only human member of the Ly-49 multigene family.